
Plants of South Eastern New South Wales



Flower and leaves. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens nursery, Canberra, ACT



Plants. Photographer Don Wood, Brogo Bush Heritage Reserve, Brogo



Line drawings. j. branchlet; k. leaves. BMMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Forest starwort

Family

Caryophyllaceae

Where found

Margins of rainforest and moist shady places in eucalypt forest, shrubland near streams, and moist gullies. Widespread. Mainly coast and ranges.

Notes

Annual or perennial herb to 0.55 m high or sprawling or scrambling, hairless or almost so, often hairy at the nodes. Sparse to medium density white hairs on new growth. Leaves opposite each other, 0.5-2.3 mm long, 1-10 mm wide, limp, margins wavy, fringed at the base, tips pointed. Flowers white, with 5 deeply cleft petals longer than the sepals. Sepals 3.5–7.5 mm long. Flowers single or in open clusters. Flowering: Spring-Summer.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Stellaria~flaccida (accessed 21 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





