# **Plants of South Eastern New South Wales**



Flower and leafy stems. Photographer Mchael Bedingfield, ACT



Back of flower. Photographer Michael Bedingfield, ACT



Plants. Photographer Michael Bedingfield, ACT



Leafy stems. Photographer Mchael Bedingfield, ACT



Line drawings (as Aster novi-belgii). plant; flower head. MMbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board,

## Common name

Michaelmas Daisy

### **Family**

Asteraceae

#### Where found

Roadsides, gardens, near habitation, gullies, and moist to swampy sites. Widespread but not common.

#### **Notes**

cm long, 3–20 mm wide; surfaces hairless or slightly hairy or bristly in lines, margins **entire** to slightly toothed, rough, bases narrowly stem-clasping. Basal parts of the plants soon leafless. Flower **heads** with few to many pale to deep mauve, purple, or bluish 'petals' each 6–15 mm long and a yellow to orange centre, 5–7 mm in diameter, with few to many **florets**. Flower heads behind the 'petals' hemispherical to bell-shaped. Flower heads in narrow leafy branched clusters. Flowers summer-autumn.

Aster novi-belgii in PlantNET.

PlantNET description (as *Aster novi-belgii*): http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Aster~novi-belgii (accessed 3 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)