# **Plants of South Eastern New South Wales**



Plants. Photographer eyew eed, Glen Oak, near New castle



Female flower and leaves. Photographer eyeweed, Glen Oak, near Newcastle



Floating male flowers. Photographer eyeweed, Glen Oak, near New castle

### Common name

Ribbonweed, Eelweed

## **Family**

Hydrocharitaceae

#### Where found

In stationary or flowing freshwater to 7m deep. Coast, ranges, tablelands, and ACT. Rarely elsewhere.

#### **Notes**

Submerged perennial aquatic herb. Leaves basal, to 300 cm long or more, 10-35 mm wide, margins toothed, tips blunt. Usually covered in brown slime and silt. Male and female flowers on separate plants. Male flowers less than 0.5 mm long, less than 1 mm in diameter. Many male flowers within a membranous **sheath**, the sheath opening to release the flowers, which float to the surface. Female flowers 1.9-4 mm long, pink or white inside, with 3 **sepals** 1-2 mm long, and 3 minute **petals**. Female flowers in clusters of 1-4. The **bracts** surrounding the female flowers translucent. Flowers Nov.—Apr..

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Vallisneria~australis (accessed 8 February, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$ 





