# **Plants of South Eastern New South Wales**



Flowers and leaves. Australian Flant Image Index, photographer Murray Fagg, Hunter Region Botanic Gardens north of Hexham



Flow ering plant. Photographer Don Wood, Biamanga National Park north of Bega

### Common name

lvy-leaved violet

## **Family**

Violaceae

## Where found

Forest, woodland, and damp places. Widespread. Rare on the tablelands north of the Hume Highway, and the Western Slopes.

#### **Notes**

Perennial herb to 0.17 m high, prostrate, stoloniferous or rhizomatous, hairless to densely hairy. Leaves in clusters along the trailing stems or forming rosettes, 0.5–3.5 cm long, 5–50 mm wide, kidney-shaped to semi-circular, margins entire to scalloped to wavy and toothed, with long stalks mostly 2–6 cm long. Flowers white with mauve to purple or violet markings, with 5 petals each 7–10 mm long, the lower petal lacking a spur, the two lateral petals often hairy within. Flowers single. Flowering: spring-summer.

Hybrids between *Viola hederacea* and *Viola sieberiana* occur with the broader leaves of the former and the smaller, more or less evenly coloured, flowers of the latter.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Viola~hederacea (accessed 8 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)