Plants of South Eastern New South Wales





Rowering plants. Australian Plant Image Index, photographer Murray Fagg, bed of Lake George, near Bungendore

Flow ering stems. Photographer Richard Hartland, Skeleton Creek, Vic?



Line drawing. e. flow ering branch. EMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Round-leafed wilsonia

Family

Convolvulaceae

Where found

In mud in coastal and subcoastal saltmarshes and brackish swamps, and in inland saline sites and the margins of salt lakes. Lake George and Lake Bathurst. Coastal north of Tuross Head.

Notes

Shrub with prostrate stems, rooting from the nodes, more or less hairy with spreading hairs. Leaves fleshy, alternating in neat rows along the stems, 0.15–0.7 cm long, 1–3 mm wide, hairy when young, more or less broad-oval with pointed tips, to oblong-spoon-shaped with blunt tips, stalkless, or with short stalks. Flowers white, cream or yellow, with a bell-shaped tube 4.5–5 mm long, and 5 lobes each 2–3 mm long. Flowers single. Flowering: mainly spring–summer.

Endangered NSW. Provisions of the NSW Biodiversity Conservation Act 2016 No 63 relating to the protection of protected plants generally also apply to plants that are a threatened species.

NSW Threatened Species profile: http://www.environment.nsw.gov.au/threatenedSpeciesApp/profile.aspx? id=10839 (accessed 8 January 2021)

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Wilsonia~rotundifolia (accessed 8 January 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:

Android edition

iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)