
Plants of South Eastern New South Wales



Flowers (spent?) and leaves. Photographer Richard Hartland, Marysville north of Melbourne, Vic



Plant. Photographer Richard Hartland, Marysville north of Melbourne, Vic



Line drawing. f. plant. K Maling, University of Sydney, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Cut-leaved xanthosia

Family

Apiaceae

Where found

Woodland, wet heath, and salong streams, and swamps. Coast, ranges, and the eastern part of the tablelands.

Notes

Perennial herb to 0.1 m high, often sprawling. Leaves mostly basal, a few alternating up the stems, 0.5-2 cm long, 15-25 mm wide, usually becoming hairless, deeply dissected into 3 lobes, the lobes often divided again, the ultimate segments linear to oval, the whole leaf almost compound. Individual flowers white or reddish, with 5 petals to about 1.5 mm long, and 5 petal-like sepals each to 1.7 mm long, mostly in clusters of 1–6 flowers. Flower clusters may look like individual flowers. Bracteoles below the flower subclusters petal-like, green or green with red tinges, 3.5–4 mm long. Flowers September–January.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Xanthosia~dissecta (accessed 8 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





