Plants of South Eastern New South Wales



Rowers and leaves. Australian Rant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Rowering branches. Photographer Don Wood, Australian National Botanic Gardens, Canberra, ACT



Shrub. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

Common name

Shapely zieria

Family

Rutaceae

Where found

Shrubland/heath on a rocky slope inland from Pambula.

Notes

Shrub to 2 m high. Stems glandular warty, and stellate-hairy (needs a hand lens or a macro app on your phone/tablet to see). Older stems less warty and becoming hairless. Leaves aromatic when rubbed, opposite each other, compound, with 3 leaflets, the central leaflet 1–3.7 cm long, 2–6 mm wide, tips more or less blunt, margins more or less toothed, and strongly curved down; upper surface warty, stellate-hairy, and light green; lower surface sparsely warty, stellate-velvety, and paler. Flowers pale pink to almost white, 6-9 mm in diameter, with 4 petals and 4 stamens. Flower clusters slightly longer than the leaves, 26-45-flowered. Bracts surrounding the flower bud clusters 1.5–6 mm long, remaining after the flowers open. Flowering: spring.

Endangered Australia. Critically Endangered NSW. Provisions of the NSW Biodiversity Conservation Act 2016 No 63 relating to the protection of protected plants generally also apply to plants that are a threatened species.

Sensitive Species list NSW.

NSW Threatened Species profile: http://www.environment.nsw.gov.au/threatenedSpeciesApp/profile.aspx? id=10855 (accessed 8 January 2021)

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Zieria~formosa (accessed 8 January 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





Creative Commons Attribution 3.0 Australia (CC BY)