

Eucalyptus kingsmillii subsp. *kingsmillii*

Classification

Eucalyptus | Symphyomyrtus | Bisectae | Destitutae | Curviptera | Xylocarpae

Nomenclature

Eucalyptus kingsmillii (Maiden) Maiden & Blakely, *Crit. Revis. Eucalyptus* 8: 43 (1929) subsp. ***kingsmillii***. 

Eucalyptus pyriformis var. *kingsmillii* Maiden, J. & Proc. Roy. Soc. New South Wales 52: 508 (1919). T: Near Mt Keith 160 miles [260 km] N of Leonora, W.A., July 1918, *W.Kingsmill s.n.*; holo: NSW; iso: CANB, K.

Description

Mallee to about 7 m tall or sometimes small **trees**. Forming a lignotuber.

Bark rough, grey and shaggy or flaky on part or all of trunk, sometimes smooth throughout with red-brown and whitish grey to grey-brown bark.

Branchlets glaucous or non-glaucous; lacking oil glands in the pith.

Juvenile growth (coppice or field seedlings to 50 cm): stems rounded in cross-section; juvenile leaves always petiolate, opposite for ca 3 nodes then alternate, ovate to lanceolate, 4–10 cm long, 1.5–4 cm wide, dull, green; new growth glaucous but the white wax not persisting as lamina expands and stems elongate.

Adult leaves alternate, petioles 0.9–2.5 cm long, lanceolate, 5.5–13 cm long; blade 1–3.6 cm wide, base tapering to petiole, concolorous, dull, blue-green to light green (new growth glaucous but wax not persisting), side-veins at an acute or wider angle to midrib, densely to very densely reticulate, intramarginal vein close to margin, oil glands few, scattered, intersectional.

Inflorescences axillary unbranched, pendulous, peduncles 1.3–4.7 cm long, buds 3 per umbel, pedicellate (pedicels 1–1.5 cm long). **Mature buds** ovoid (2.2–4 cm long, 1.5–2.5 cm wide), with ca 8 longitudinal ribs, glaucous or non-glaucous, scar present, operculum beaked or conical (1.5–2.8 cm long), outer stamens erect, inner inflexed, or all stamens more or less obliquely arranged, anthers oblong, versatile, sub-dorsifixed, dehiscing by lateral slits, style long, stigma blunt, locules 3, 4 or 5, the placentae each with 6 vertical ovule rows (or with 6 rows increasing to 8 rows basally). Flowers yellow sometimes with basal tinge of pink to filaments.

Fruit down-turned, pedicellate (pedicels 0.5–3 cm long), hemispherical to obconical, 0.8–2 cm long, 1.5–4 cm wide (including the often prominent ribs), rarely glaucous, disc raised and concave, valves 3, 4 or 5, exerted.

Seeds greyish brown, 2–3.5 cm long, more or less flattened-pyramidal, dorsal surface shallowly reticulate, sides ridged, encircling marginal flange usually present, hilum terminal.

Cultivated seedlings (measured at ca node 10): cotyledons Y-shaped (bisected); stems rounded in cross-section; leaves always petiolate, opposite for 4 to 6 nodes then alternate, ovate-lanceolate, 4.5–10 cm long, 1.8–4 cm wide, dull, greyish green to green.

Flowering Time


Flowering has been recorded in March, April, May, August and September.

Notes

Eucalyptus kingsmillii is a mallee species widespread in arid central and south-eastern Western Australia, extending just into South Australia near Serpentine Lakes. The bark is rough and loose over the lower part of the stems and usually ribbony above. The adult leaves are dull, blue-green to light green.

Eucalyptus kingsmillii belongs in *Eucalyptus* subgenus *Symphyomyrtus* section *Bisectae* subsection *Destitutae* because buds have two opercula, cotyledons are Y-shaped and branchlets lack oil glands in the pith. Within this subsection *E. kingsmillii* is in series *Curviptera*, one of about 30 closely related species and subspecies which are further characterised by having large buds in umbels of one, three or seven, staminal filaments erect or oblique (rarely inflexed) in bud, and large fruit usually with an ascending disc and exerted valves. Several of these species are noted for their large-leaved crowns made up of juvenile leaves.

E. kingsmillii is closely related to *E. youngiana* and two more northerly species, *E. sessilis* and *E. pachyphylla*. All are desert-dwelling species with prominently ribbed buds and fruit. *E. pachyphylla* has buds in threes on short erect stalks and *E. sessilis* has buds also on



short erect stalks but mostly in sevens. *E. youngiana* and *E. kingsmillii* both have buds in threes on downturned peduncles. The fruit of *E. kingsmillii* (to 4.5 cm wide) are smaller than those of *E. youngiana* (to 7.2 cm wide). The flowers of *E. youngiana* are brilliant red, pink or bright yellow while *E. kingsmillii* subsp. *kingsmillii* has pale yellow flowers sometimes suffused basally with pink, and *E. kingsmillii* subsp. *alaticissima* has pink or red flowers, rarely creamy yellow. The desert dwelling mallee species *E. oxymitra* differs in having smaller un-ribbed buds and fruit.

There are two subspecies:

E. kingsmillii* subsp. *kingsmillii

This is the common widespread form which is endemic to Western Australia, occurring south from the Hamersley Range to Leonora, east to the southern Gibson Desert and Warburton, on mountain-tops and other elevated rocky sites, sand dunes, sandplain. Branchlets, buds and fruit are rarely glaucous.

E. kingsmillii* subsp. *alaticissima

This form is from the Great Victoria Desert extending eastwards from Warburton (WA) into the Serpentine Lakes area of South Australia on red sand dunes and sandplain. It differs by having glaucous branchlets, more attenuate opercula and much stronger ribbing of the fruits.

Origin of Name

Eucalyptus kingsmillii : after Sir Walter Kingsmill (1864–1935). Walter Kingsmill was a state politician of Western Australia and held the seat of Pilbara from 1897–1903. From an early stage in his political career he held ministerial rank, frequently becoming Acting-Premier of Western Australia. In 1922 he was elected to the Australian Senate and held that position until his death. He was President of the Senate from 1929 to 1932.

subsp. *alaticissima*: Latin *alatus* - winged, in the superlative, referring to the fruit

Copyright © CANBR 2020, all rights reserved.



Web edition hosted at <https://apps.lucidcentral.org/euclid>