

# *Crepidomanes barnardianum*

## Family:

Hymenophyllaceae

## Botanical name

*Crepidomanes barnardianum* (F.M.Bailey) Tindale

Link to Australian Plant Name Index for publication details and synonyms:

<https://id.biodiversity.org.au/name/apni/84387>

## Description

Rhizome to 0.25 mm diam., widely creeping; hairs dark reddish brown, extending to stipe margins. Fronds 0.7–4 cm long. Stipe 1–10 mm long, winged to the base. Lamina ovate to ± palmate, 1–2 (–3)-pinnatifid, 0.7–3 cm long, 0.4–3.4 cm wide, glabrous; ultimate segments less than 1 mm wide, acute or obtuse; submarginal false vein present, mainly continuous; accessory false veins few. Sori clustered towards the frond apex; involucre cylindrical, tapering towards the base, 1–1.8 mm long, 0.7–1.1 mm diam.; mouth ± dilated but not bilabiate; receptacles exerted to c. 3 mm beyond the mouth.

## Distribution

Endemic to NE QLD between Iron Range and the Whitsundays.

## Habit and habitat

Lithophytic or epiphytic, usually in shaded strata of mesic lowland to upper montane vine forest.

## Citation of Australian Tropical Ferns and Lycophytes

Field AR, Quinn CJ, Zich FA (2022) Australian Tropical Ferns and Lycophytes. [apps.lucidcentral.org/fern/text/intro/index.htm](https://apps.lucidcentral.org/fern/text/intro/index.htm) (accessed online INSERT DATE).

Field AR, Quinn CJ, Zich FA (2022) '*Platyserium superbum*', in Australian Tropical Ferns and Lycophytes. [apps.lucidcentral.org/fern/text/entities/platyserium\\_superbum.htm](https://apps.lucidcentral.org/fern/text/entities/platyserium_superbum.htm) (accessed online INSERT DATE).



Close up of frond showing sori.

© G. Sankowsky



Close up of frond showing sori.

© CSIRO



Habit. © G. Sankowsky



Habit. © G. Sankowsky



Habit. © CSIRO

