

# *Oreogrammitis wurunuran*

**Family:**

Polypodiaceae

**Botanical name**

*Oreogrammitis wurunuran* (Parris) Parris

Link to Australian Plant Name Index for publication details and synonyms:

<https://id.biodiversity.org.au/name/apni/215199>

**Description**

Rhizome erect. Fronds c. 0.2 mm apart. Stipe with medium to dark red-brown simple eglandular hairs 0.3–1.3 mm long. Lamina (1.8–) 2.8–6.6 (–12.5) cm long, (3–) 4–6 (–7) mm wide, thinly coriaceous to coriaceous; hairs dark red-brown, simple, 0.3–1.7 (–3) mm long, eglandular, on margin and both surfaces of lamina and midvein; veins usually not visible in transmitted light, without hydathodes. Sori rounded to elliptic, (1–) 6–19 (–27) pairs in middle to apical one-fifth to five-sixths of lamina. Sporangia setose; thickened annulus cells 9–12 (–15). Spores (23–) 26–33 (–39) µm diam.

**Distribution**

Endemic to NE QLD between Mt Finnegan and Eungella.


**Habit and habitat**

Epiphytic and lithophytic in mid- to upper-montane rainforest.

**Citation of Australian Tropical Ferns and Lycophytes**

Field AR, Quinn CJ, Zich FA (2022) Australian Tropical Ferns and Lycophytes. [apps.lucidcentral.org/fern/text/intro/index.htm](https://apps.lucidcentral.org/fern/text/intro/index.htm) (accessed online INSERT DATE).

Field AR, Quinn CJ, Zich FA (2022) '*Platyserium superbum*', in Australian Tropical Ferns and Lycophytes. [apps.lucidcentral.org/fern/text/entities/platyserium\\_superbum.htm](https://apps.lucidcentral.org/fern/text/entities/platyserium_superbum.htm) (accessed online INSERT DATE).



Close up of frond showing sori.  
© G. Sankowsky



Close up of frond showing sori.  
© CSIRO



Frond showing upper (adaxial) side. © G. Sankowsky



Habit. © G. Sankowsky



Habit. © CSIRO



Habit. © G. Sankowsky

