# Schizaea dichotoma

Family: Schizeaceae

#### Botanical name

Schizaea dichotoma (L.) Sm.

Link to Australian Plant Name Index for publication details and synonyms: https://id.biodiversity.org.au/name/apni/116177

# Description

Rhizome short-creeping, densely clothed with glossy dark brown hairs. Fronds clustered, erect, (3.5–) 10–50 cm long, scabrid; unbranched portion 1–2 mm wide, subterete, flattened on 1 side, brown near base, clothed with long brown hairs, green above; hairs scattered; sterile frond shorter than sporogenous frond, dividing above middle into a dichotomously branched, flabellate portion, with 20–180 flat asperous acute branchlets; sporogenous frond similar, but branching more open; sporogenous heads 7–15 mm long, 1–3 times longer than broad; segments 3–9 mm long, scabrid; margins with numerous long hairs. Sporangia mixed with hairs.

# Distribution

Widespread across tropical and subtropical coastal Australia.

# Habit and habitat

Terrestrial, usually in vine forest, particularly beach scrubs and sandy areas adjacent watercourses. Also in rocky areas in open mid-montane vine forest.

#### Cultivation

Considered difficult to propagate and establish.

#### Citation of Australian Tropical Ferns and Lycophytes

Field AR, Quinn CJ, Zich FA (2022) Australian Tropical Ferns and Lycophytes. apps.lucidcentral.org/fern/text/intro/index.htm (accessed online INSERT DATE).

Field AR, Quinn CJ, Zich FA (2022) 'Platycerium superburn', in Australian Tropical Ferns and Lycophytes. apps.lucidcentral.org/fern/text/entities/platycerium\_superburn.htm (accessed online INSERT DATE).

Copyright © Australian Tropical Herbarium 2022, all rights reserved.



Frond. © A.R. Field



Frond. © G. Sankowsky



Close up of frond showing sori. © G. Sankowsky



Close up of frond showing sori. © G. Sankowsky



Habit. © CSIRO



Habit. © G. Sankowsky



Frond. © A.R. Field

14



Web edition hosted at https://apps.lucidcentral.org/ferns