

Plants of South Eastern New South Wales



Flower heads, buds, and 'leaves'. Photographer Jackie Miles



Flower heads, pods, and 'leaves'. Photographer Don Wood, west of Nowra



'Leaf'. Photographer Jackie Miles



Shrub. Photographer John Tann, near Sydney

Common name

Two-veined hickory

Family

Fabaceae

Where found

Forest and moist sites. Coast and ranges north from the Moruya district.

Notes

Shrub or tree to 15 m tall. Fleshy seed stalks/**arils**. Bark smooth or rough, more or less flaky and fissured. Branchlets angled towards the tips, hairless. 'Leaves' **alternating** up the stems, 5-16 cm long, 10-50 mm wide, straight to curved, surfaces hairless, tips pointed, 1 **marginal gland** near the base. Flower **heads** globular, white to pale yellow, 15-50 flowered (easiest seen in late buds), 5-10 mm in diameter, in clusters of 3-12 flower heads. Flowers Winter to Spring.

Family was Mimosaceae.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia-binervata> (accessed 3 April, 2021)

World Wide Wattle line drawings, photos, and description: <http://worldwidewattle.com/imagegallery/image.php?p=0&l=b&id=23571&o=1>

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Creative Commons Attribution 3.0 Australia (CC BY)



Android edition



iOS edition