Plants of South Eastern New South Wales



Flowering stem Australian Plant Image Index, photographer Murray Fagg, Ingalba Nature Reserve west of Temora



Flowering stems. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Shrubs. Australian Flant Image Index, photographer Mırray Fagg, İngalba Nature Reserve Griffith to Termra



Line drawing. f. flowering branch. MMbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Bent-leaf wattle, Small winter wattle

Family

Fabaceae

Where found

Dry forest, woodland, and shrubland. Western Slopes. Doubtfully naturalised in the ACT.

Notes

Shrub to 1.5 m tall, spreading or sprawling. Fleshy seed stalks/arils. Branchlets angled at the tips, hairy with minute hairs, and scurfy between resinous longitudinal ridges. 'Leaves' alternating up the stems, 1-2.5m long, 0.7-2 mm wide, straight or slightly curved, surfaces hairless, veins and margins resinous, midvein prominent, closer to the upper margin and often joining it towards the tip, tips blunt with a mucro. 1 marginal gland near the base, the margin usually notched at the gland. Flower heads yellow, globular, 4-6 mm in diameter, 3-10 flowered (easiest seen in late buds), usually in pairs. Flowering: usually June—September.

Family was Mimosaceae.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia~flexifolia (accessed 27 April 2021)

World Wide Wattle line drawings, photos, and description: http://www.worldwidewattle.com/imagegallery/image.php?
p=0&l=f&id=23652&o=1

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)