
Plants of South Eastern New South Wales



Flowering stems and leaves. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Pods and leaves. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra

Common name

A wattle

Family

Fabaceae

Where found

Dry forest and woodland, often on stony ridges. Ranges and the eastern edge of the tablelands north of inland from Jervis Bay.

Notes

Shrub to 4 m tall. Bark smooth. Fleshy seed stalks/arils. Branchlets angled to cylindrical with ridges, hairy with minute hairs, or hairless. Leaves alternating up the stems, compound, rachis 1-7.5 cm long, hairy to hairless, a small round more or less jugary gland present at the base of or just below each pair of pinnae except at the basal pinnae, rarely missing from the others, interjugary glands absent. 4-22 pinnae, 0.5-4 cm long, each with 8-42 leaflets each 0.25-0.6 cm long, 0.8-2 mm wide, hairless, or hairy with minute hairs mainly on the margins. Flower heads cream to yellow, globular, 8-15 flowered (easiest seen in late buds), 5-8 mm in diameter, in elongated clusters of 6-25 flower heads. Flowers Winter to Spring, occasionally late Summer.

Family was Mimosaceae.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia~jonesii (accessed 27 April 2021)

World Wide Wattle line drawings and description: http://www.worldwidewattle.com/imagegallery/image.php?
p=0&l=j&id=23693&o=1

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)