

Plants of South Eastern New South Wales




Flowering stems. Australian Plant Image Index, photographer Murray Fagg, Wyperfeld National Park, Vic



Stems with pods. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Shrub. Australian Plant Image Index, photographer Murray Fagg, Wyperfeld National Park, Vic



Line drawings. c. flowering branch with stem detail. MIMbir, M., National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Mallee wattle

Family

Fabaceae

Where found

Dry **forest**, **woodland**, and **shrubland**. Western Slopes.

Notes

Shrub or tree 0.5–4 m tall. Fleshy seed stalks/**arils**. Bark fissured. Branchlets angled towards the tips, resinous, densely hairy to almost hairless. 'Leaves' **alternating** up the stems, 1-4.5 cm long, 1.5-7 mm wide, hairless or sparsely hairy, shiny and/or resinous, tips rounded with a very short point. Flower **heads** yellow, globular, 4-7 mm in diameter, 20-30 flowered (easiest seen in late buds), single or paired. Flowering: usually August–November. Pods pale fawn-**woolly** to hairy.

Family was Mimosaceae.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvi=sp&name=Acacia-montana> (accessed 27 April 2021)

World Wide Wattle line drawings, photos, and description: <http://www.worldwidewattle.com/imagegallery/image.php?p=0&l=m&id=23745&o=1>

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Creative Commons Attribution 3.0 Australia (CC BY)



Android edition



iOS edition