

Plants of South Eastern New South Wales



Flowering stems and 'leaves'. Photographer Don Wood, Australian National Botanic Gardens, Canberra




Flowering stems. Photographer Jackie Mles



Leafy stems with young pods. Photographer Jackie Mles



Shrub, photographer Jackie Mles, near Mogo south of Batemans Bay



Line drawing. h. flowering branch with 'leaf' detail. M Mbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Net-veined wattle, Stiff-leaf wattle

Family

Fabaceae

Where found

Dry **forest** and **heath** on hillsides and rocky outcrops, sometimes near streams. Coast and ranges. Tablelands north from Goulburn.

Notes

Tree or **shrub** to 5.5 m tall. Fleshy seed stalks/**arils**. Bark smooth, somewhat fibrous, or finely fissured. Branchlets more or less

cylindrical, hairless, more or less resinous, at first covered with resin hairs towards the tips. 'Leaves' **alternating** up the stems, 5-19.5 cm long, 3-16 mm wide, leathery, surfaces hairless, **midvein** prominent, tips blunt or pointed, often with a **mucro**. Flower **heads** yellow, oval to cylindrical, 10-30 mm long, single or in pairs. Flowers Winter to Summer.

Family was Mimosaceae.

Vulnerable Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia~subtilinervis> (accessed 28 December, 2020)

World Wide Wattle line drawings and description: <http://www.worldwidewattle.com/imagegallery/image.php?p=0&l=s&id=23842&o=1>

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition