

Plants of South Eastern New South Wales




Flowering branches. Australian Plant Image Index, photographer Murray Fagg, Alleena to Tallirba, NSW



Leafy stems with open pods. Photographer Lorraine Phelan, Lake Hindmarsh, Vic



Shrub. Australian Plant Image Index, photographer Murray Fagg, Alleena to Tallirba, NSW



Line drawing. e. flowering branch with 'leaf' detail. M Moir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Three-nerved Wattle, Three-veined Wattle, Green Wattle, Hindmarsh Wattle

Family

Fabaceae

Where found

Mallee, sometimes near water. Western Slopes, mainly in the West Wyalong-Condobolin-Forbes area.

Notes

Shrub or tree to 5 m tall, sometimes almost **prostrate**. Fleshy seed stalks/**arils**. Bark smooth, grey. Branchlets more or less cylindrical, resin-ribbed at the extremities, densely whitish- or greyish-hairy with minute hairs, becoming hairless with age. 'Leaves' **alternating** up the stems, 3–8 cm long, 3–11 mm wide, somewhat **glaucous**, hairless or hairy with fine hairs, 3–7 main longitudinal **veins**, 3 often slightly more prominent, tips blunt with a **mucro**. Flower **heads** pale yellow to yellow, 3–5 mm in diameter, 12 to about 25 flowered (easiest seen in late buds), globular, in 3–8 flowered clusters. Flowering: August–October. **Aril** folded 1–3 times.

Family was Mimosaceae.

Vulnerable Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswf1&lvi=sp&name=Acacia-trineura> (accessed 28 April 2021)

World Wide Wattle line drawings, photos, and description: <http://worldwidewattle.com/imagegallery/image.php?p=0&l=t&id=23854&o=1>

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)