
# **Plants of South Eastern New South Wales**



Flow ering plant. Photographer Chris Clarke, Lake Mountain, Vic



Flow ering plants. Photographer Russell Best, near Gisborne, Vic



Line drawings (as *Brachyscome cardiocarpa*). a. flowering branch; plant; seed (front view). Goria Thontinson, Daisy Study Group, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

### Common name

Swamp daisy

### **Family**

Asteraceae

## Where found

Swampy areas. Ranges south from Deua National Park. One record from near Yaouk, north of Adaminaby.

#### **Notes**

Perennial herb to 0.45 m tall, mainly hairless. Leaves mainly basal, 4-30 cm long, 1-3 mm wide, mainly hairless but at least the lower leaves with long, coarse, brownish hairs at the base, margins usually entire, very rarely with several small linear lobes. A few narrow stem leaves, which reduce in length upwards towards the flower head. Flower heads behind the 'petals' almost cone shaped, with 54-77 white or mauve 'petals' each 10-16.2 mm long, and yellow centres about 8-15 mm in diameter. Bracts appressed to the flower heads 25–40. Flowering: mostly Jun.–Feb.

Was Brachyscome cardiocarpa

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Allittia~cardiocarpa (accessed 2 January, 2021)

Description mainly based on Short, P.S., (2014) A taxonomic review of *Brachyscome* Cass. s.lat. (Asteraceae: Astereae), including description of a new genus, *Roebuckia*, new species and new infraspecific taxa. *Journal of the Adelaide Botanic Gardens*. 28: 55-57

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)