Plants of South Eastern New South Wales



Flowers and leaves. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ering branches. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

Common name

None

Family

Rutaceae

Where found

Wet **forest** in moist gullies. Ranges and eastern edge of the tablelands north of Kangaroo Valley. Coastal north from Sydney. Doubtfully **naturalised** in the ACT.

Notes

Shrub to 2 m high. Branches white to brown stellate hairy (needs a hand lens or a macro app on your phone/tablet to see). Leaves aromatic when rubbed, alternating up the stems, 3–9 cm long, 8–30 mm wide, tips blunt to pointed, upper surface hairless and smooth, lower surface white to brownish stellate-hairy. Flowers white to cream, brown stellate-hairy outside, with 5 petals each 5–6 mm long, usually in clusters of several flowers. Flowering: spring.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Asterolasia~correifolia (accessed 4 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)