Plants of South Eastern New South Wales



Closeup of flow er head. Australian Plant Image Index, photographer A Lyne Australian National Botanic Gardens, Canberra, ACT

Common name

A Banksia

Family

Proteaceae

Where found

Dry forest and woodland in rocky areas. Ranges north from Nattai National Park.

Notes

Shrub to 4 m high. Bark smooth, thin, grey. Branchlets more or less striate, rusty-hairy when young. Leaves whorled, 4–12 cm long, 10-40 mm wide, margins usually toothed or lobed, lower surface white-hairy, midrib and lateral veins with more or less persistent rusty hairs, tips blunt and often mucronate. Flower heads 70-190 mm long, of many flowers. Individual flowers yellowish green to bluish grey in bud, yellow in open flowers, with 4 'petals' each 20-25 mm long, splitting to the base when the flowers are fully open. Style slightly curved, pale yellow, remaining on mature cones. Flowers March-June. Seeds with one wing.

Hybrids with Banksia marginata have been recorded.

PlantNET description (as Banksia penicillata: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Banksia~penicillata (accessed 17 April 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:



Android edition iOS edition - 1

Creative Commons Attribution 3.0 Australia (CC BY)