Plants of South Eastern New South Wales



Flowers. Photographer Teun Spaans, unknown place



Flow ering stem. Photographer S Drozd Lund, Denmark



Fruit and leaves. Photographer Algirdas, unknown place



Shrub. Photographer Teun Spaans, unknown place



Line drawings. USDA-NRCS PLANTS Database / Britton, NL., and A. Brown. 1913. An illustrated flora of the northern United States, Canada and the British Possessions. Vol. 2: 127.

Common name

Common Barberry, Barberry

Family

Berberidaceae

Where found

Weed. Southern Highlands mainly between Mittagong and Bundanoon. Kosciuszko National Park. Occasionally elsewhere.

Notes

Introduced deciduous shrub to 2.5 m high. Leaves on the main stems reduced to spines, usually deeply divided into 3 spines, each 10–25 mm long. Fruit fleshy. Stems hairless. Leaves clustered on short shoots, 2–6 cm long, 8–25 mm wide, margins toothed with numerous bristly teeth. Flowers yellow, about 5 mm in diameter, with 6-12 petals in 2 or 3 whorls. Flowers in many-flowered clusters 5–7 cm long. Fruit bright red, oblong to oval, 9–12 mm long, edible.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)