Plants of South Eastern New South Wales



Flowering branch. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flowering branches. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Shrub. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens,

Common name

Native Rose, Rose Boronia

Family

Rutaceae

Where found

Moist heath in sandy situations. Coast, and occasionally ranges, north of Berry.

Notes

Shrub to 1.5 m high. Branchlets with 2 grooves separated by the ridges formed by the leaf bases continuing down the stems, and hairless or rarely sparsely hairy with simple hairs. Leaves strongly aromatic when rubbed, crowded, 0.7–1.8 cm long, 5–9 mm wide, hairless, thick, often red-tinged, margins finely toothed, tips pointed. Flowers bright pink, with 4 petals each 7–11 mm long. Petals overlapping each other in bud, falling early. 8 stamens. Flowers in 1–4-flowered clusters. Flowering: August–October.

Protected NSW.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Boronia~serrulata (accessed 5 January, 2020)

Author: Betty Wood.

 $This \, identification \, key \, and \, fact \, sheets \, are \, available \, as \, a \, \, free \, mobile \, application: \, \\$





