Plants of South Eastern New South Wales



Flow ering stems. Photographer Don Wood, Morton National Park



Shrubs. Australian Plant Image Index, photographer lan Telford, Mt Budawang east of Braidwood

Common name

A bossiaea

Family

Fabaceae

Where found

Dry forest and heath, on exposed rocky sites, and in moist habitats. Coast, ranges, and the eastern part of the tablelands, mainly between the Kings Highway and the Illawarra Highway. Occasionally elsewhere.

Notes

Shrub, prostrate or to 4 m tall. Stems mildly compressed at first, becoming cylindrical, hairy with white hairs, becoming hairless. Leaves opposite each other, 1-3.5 cm long, 2-7 mm wide, upper surface hairless, lower surface sparsely hairy to hairless, paler than the upper surface, tips with a short point. Flowers usually 10-12 mm long, pea-shaped with 5 petals, 2 joined together to form the keel. Front of standard petal yellow with a red flare, back generally red, wings light purplish brown with red streaks,, keel red to dark brown. Flowers Spring.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Bossiaea~kiamensis (accessed 5 January, 2020)

Description partly based on Thompson, I.R., (2012) A revision of eastern Australian *Bossiaea* (Fabaceae: Bossiaeae). *Muelleria* 30(2): 127-128

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)