
# **Plants of South Eastern New South Wales**



Flowering plants. Australian Flant Image Index, photographer Colin Totterdell, Kosciuszko National Park



Flow ering plant. Photographer Don Wood, Namadgi National Park, ACT



Line drawings. b. flowering branch; seed (front and side view). Gloria Thontinson, Daisy Study Group, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

#### Common name

**Tufted Daisy** 

### **Family**

Asteraceae

## Where found

Forest, woodland, heath, alpine herbfields and grasslands, and swamp margins. Kosciuszko National Park, the mountains to the north, ACT, tablelands, and the ranges.

#### **Notes**

Perennial herb to 0.48 m high. Flower stalks hairless or hairy with few to many glandular hairs, usually with 1-2 (rarely more) small leaf-like bracts. Leaves basal, 2.5-24 cm long, 3-15 mm wide, hairless or rarely with some glandular hairs, margins entire, tips pointed. Plant with the remains of old leaves persisting at the base of the flower stalk(s). Flower heads single, 15-30 mm in diameter overall, with 45-90 white to mauve 'petals' 6-10.7 mm long, and yellow centres 6-11 mm in diameter. Bracts appressed to the flower heads 14-25. Flowering: Oct-May.

Definite identification of most species of Brachyscome depends on microscopic examination of the seeds.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Brachyscome~scapigera (accessed 6 January, 2021)

Description partly based on: Short, P.S. (2014), A taxonomic review of *Brachyscome* Cass. s.lat. (Asteraceae: Astereae), including description of a new genus, *Roebuckia*, new species and new infraspecific taxa. *Journal of the Adelaide Botanic* Gardens 28: 158

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)