
Plants of South Eastern New South Wales



Flow ering plant. Photographer Don Wood, Namadgi National Park, ACT



Rowering plant. Photographer Don Wood, Ben Boyd National Park south of Eden



Line drawings (as *Brachyscome spathulata* subsp. *spathulata*). e. plant; seed (front view). Gloria Thorninson, Daisy Study Group, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Coarse daisy, Spoon-leaved daisy

Family

Asteraceae

Where found

Forest, woodland, heath, very low shrubland in alpine feldmark, and grassy areas and herbfields. Widespread. No records from the Sydney area.

Notes

Perennial stoloniferous herb to 0.6 m high. Scapes sticky hairy to more or less hairless. Most leaves in a basal rosette. Basal leaves spoon-shaped, 1.7-12 cm long, 6-30 mm wide, more or less hairless, margins mostly with shallow teeth or lobes. Stem leaves alternating up the stems, 0.5–5 cm long, 1–30 mm wide, woolly, margins finely toothed, becoming smaller in size up the stem. Flower heads single, with 40-60 pale blue to mauve or rarely white 'petals' 5–16 mm long, and yellow centres 10–20 mm in diameter. Bracts appressed to the flower heads about 20-30. Flowering: Sept.–June. 'Seeds' flat.

All native plants on unleased land in the ACT are protected.

Definite identification of most species of Brachyscome depends on microscopic examination of the seeds.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Brachyscome~spathulata (accessed 6 January, 2021)

Description partly based on: Short, P.S. (2014), A taxonomic review of *Brachyscome* Cass. s.lat. (Asteraceae: Asteraee), including description of a new genus, *Roebuckia*, new species and new infraspecific taxa. *Journal of the Adelaide Botanic Gardens* 28: 39-41

Author: Betty Wood.

6 ۲ This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)