Plants of South Eastern New South Wales



Flow ers and leaves. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens



Flowering plant. Photographer Don Wood, west of Nowra

Common name

A daisy

Family

Asteraceae

Where found

Dry forest, woodland, and shrubland. North of the Hume and Illawarra Highways. Occasionally farther south.

Notes

Perennial herb with stems to 0.35 m long, sprawling to weakly erect, hairless. Rhizomatous. Leaves basal and alternating up the stems, 0.7-7.5 cm long, 0.5-19 mm wide, hairless or with very occasional glandular hairs, margins usually entire, occasionally with with 1-5 (mostly 3) lobes at the tips, the middle lobe much larger than the other two. Flower heads with about 21-30 or more 'petals' about 7-7.5 mm long, pink, mauve, or blue to purple, and yellow centres about 6-7 mm in diameter. Bracts appressed to the flower heads in one row. Flowers throughout the year.

Was Brachyscome angustifolia, Brachyscome linearifolia.

Definite identification of most species of *Brachyscome* depends on microscopic examination of the seeds.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Brachyscome~triloba (accessed 6 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)