


Plants of South Eastern New South Wales



Flowering plants. Australian Plant Image Index, photographer Murray Fagg, Nadgigomar Nature Reserve north of Braidwood



Leaves. Photographer Richard Hartland, Terrick Terrick, Vic



Line drawings. j. plants; seed. E Mayfield, National Herbarium of Victoria, ©2021 Royal Botanic Gardens Board

Common name

Cut-leaved burr-daisy, Burr Daisy

Family

Asteraceae

Where found

Open **grassland**, mostly in depressions or damp areas. ACT. Tablelands and the western edge of the ranges east and north of the ACT, mainly south of the Hume Highway.

Notes

Perennial herb to 0.2 m high, **stoloniferous**, forming colonies, hairless or with a few small scattered hairs. 'Seeds' flattened, with rigid **wings** on the margins, and 7–14 unequal spines to 2 mm long, with minute barbs at the tips. 'Seeds' stick to clothing. Leaves hairless, basal, in a **rosette**, 3–12 cm long, 6–20 mm wide, **compound** or deeply **dissected**, with narrow lobes/leaflets, hairless, with long stalks. Flower **heads** 10–20 mm in diameter, single, with 40–70 white '**petals**' 4–7 mm long, and yellow centres. Flowering: Sept.–Feb.

Definite identification of all species of *Calotis* is made by examining the seeds.

All native plants on unleased land in the ACT are protected.

Listed in the Flora and Fauna Guarantee Act, Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Calotis-anthemoides> (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)