


# Plants of South Eastern New South Wales



Flower, buds, and leaves. Photographer Richard Hartland, Noojee, Vic?



Flower bud showing bracteole, and leaf. Photographer Bill Campbell, New Zealand

## Common name

Forest Bindweed

## Family

Convolvulaceae

## Where found

**Forest**, moist gullies, and stream banks. Coast, ranges, and the eastern part of the tablelands.

## Notes

**Perennial** herb with twining stems. Hairless. Leaves **alternating** along the stems, 3–10 cm long, 15–80 mm wide, arrow **head** shaped, basal lobes toothed or shallowly lobed, tips pointed. Flowers white to pale pink or mauve, about 20 mm long, trumpet-shaped, with 5 lobes. **Stigma** with 2 lobes. **Sepals** free from each other. **Bracteoles** oval to round, tips rounded to notched, 6–12 mm long, equal to or longer than the **calyx**. Flowers single. Flowering: spring–summer.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Calystegia-marginata> (accessed 7 January, 2021)

*Author: Betty Wood.*



*Creative Commons Attribution 3.0 Australia (CC BY)*

*This identification key and fact sheets are available as a free mobile application:*

