Plants of South Eastern New South Wales



Flowers. Australian Flant Image Index, photographer Murray Fagg, Barrington Tops National Park



Line drawing. d. flowering plant. MMbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Spreading bitter-cress

Family

Brassicaceae

Where found

Grassy areas in alpine and sub-alpine areas. Kosciuszko National Park, the mountains to the north, and ACT.

Notes

Perennial herb with stems growing along the ground, rooting at the nodes and producing erect unbranched flowering stems to 0.3 m high. Basal leaves single along the stems, sometimes a few clustered at the base of the flowering stem, to 15 cm long, sometimes compound, with 2 or 4 lateral leaflets and a terminal leaflet. Stem leaves alternating up the stems, to 6 cm long, deeply dissected or compound. Flowers with 4 petals, white or pink on both sides, or pink on the outside, 6–11 mm long. 6 stamens. Flowers in many-flowered clusters. Flowers late spring to summer.

Vulnerable Vic.

PlantNET description with line drawing and photos of dried specimens: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Cardamine~astoniae (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)