Plants of South Eastern New South Wales



Flow ering plant. Photographer lan McMaster, Kosciuszko National Park

Common name

A Bitter-cress

Family

Brassicaceae

Where found

Moist boulder screes and stream and lake margins. Kosciuszko National Park and ACT. Western edge of the ranges south of Bombala.

Notes

Rhizomatous perennial herb to 0.3 m high, sprawling, forming dense swards up to 1 m or more in diameter. Stems hairless, with conspicuous leaf scars towards the base. Basal leaves forming a **rosette** distinctly above ground level, to 25 cm long, **compound**, with 3-7 leaflets. Stem leaves 0–4, **alternating** up the stems, similar to the basal leaves but becoming shorter up the stem, the leaflets becoming narrower. Flowers with 4 white **petals** each about 8–12 mm long, and 6 **stamens**. Flowers in few- to many-flowered clusters. Flowers late spring to early autumn.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Cardamine~robusta (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)