
Plants of South Eastern New South Wales



Flowering plants. Australian Flant Image Index, photographer Murray Fagg, Kosciuszko National Park



Line drawings. e. leaf; leaf margin detail. MMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Sticky Billy-buttons

Family

Asteraceae

Where found

At high altitude. Heath, grassland, alpine herbfields, margins of alpine feldmark, and rocky outcrops. Kosciuszko National Park, the mountains to the north, and ACT.

Notes

Herb to 0.4 m high, sticky. Flower stalks straw-coloured, brown to red-brown, sticky, sometimes with appressed silky hairs. Leaves smell like mouldy oranges or citronella when rubbed. Leaves mostly basal, 6.5-18 cm long, 8-20 mm wide, upper surface dark green with dense glandular hairs, sometimes with loose appressed woolly hairs, particularly on the midrib, lower surface densely appressed woolly tomentose with short glandular hairs intermixed, tips pointed. Flower head usually yellow, sometimes pale yellow or orange, about 15-35 mm in diameter, with about 50–100 partial heads. Partial heads with 5-10 florets with 0 petals. Flower head hemispherical to globular, single.

Protected NSW.

All native plants on unleased land in the ACT are protected.

Rare Vic.

It appears to form hybrids with Craspedia maxgrayi where their ranges overlap.

VICFLORA description: https://vicflora.rbg.vic.gov.au/flora/taxon/df0af0fc-2edd-44b4-8f16-cf50cc95d0a8 (accessed 9 February 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)