


Plants of South Eastern New South Wales



Flowering plant. Photographer Richard Hartland, Anglesea, Vic



Plant with flower buds. Photographer MFdeS, Tas



Line drawing. f. leaf and stipules. R Barley, Royal Botanic Gardens Melbourne, © 2021 Royal Botanic Gardens Board

Common name

Pimpernel sundew, Scarlet sundew

Family

Droseraceae

Where found

Moist places and swamps in dry **forest**, **woodland**, **heath**, and **grassland**. Coast and ranges north from west of Jervis Bay, and south of Merimbula. Western Slopes.

Notes

Annual herb to 0.1 m high, **rhizomatous**. Leaves basal, in a **rosette**; 8–20 mm in diameter, almost round, very concave in cross section, abruptly narrowed to the leaf stalk, upper surface covered with **glandular** hairs that trap insects. **Stipules** more or less **entire**, fused to the base of the leaf stalk, 1–2.5 mm long. **Petals** orange or red, 4–5 mm long. Flowers in 1–8 clusters, each about 7–many-flowered, glandular; individual stalks bending down after flowering. Flowering: winter to spring.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvi=sp&name=Drosera-glanduligera> (accessed 12 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)