
Plants of South Eastern New South Wales



Flowering stems. Photographer Richard Hartland, Bogong Hgh Plains, Vic



Flow ering plant. Photographer Don Wood, Namadgi National Park



Line drawings. d. plant; e. seed. E.Mayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Bald-seeded willow-herb

Family

Onagraceae

Where found

Moist places in **grassland** or in temporary shallow pools. At higher altitudes. Kosciuszko National Park, the mountains to the north, and the ACT. Rarely elsewhere.

Notes

Mat-forming **perennial** herb, to 0.15 m high where vegetation is dense. Lower branches rooting in mud. Stems sparsely to densely covered with wavy hairs. Leaves opposite each other on the lower stems, **alternating** along the stems near the flowers, 0.5–2.5 cm long, 2–9 mm wide, margins with sparse fine teeth, and often wavy. Flowers with 4 pale purplish pink **petals** 3.5–7 mm long, the inside of the **floral tube** hairless or with a few long hairs. Seed cases 0.8–2.5 cm long, very sparsely hairy, on stalks 0–3 mm long.

Definite identification of Epilobium species requires microscopic examination of the seeds.

All native plants on unleased land in the ACT are protected.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Epilobium~curtisiae (accessed 1 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





