

# Plants of South Eastern New South Wales



Flowers and leaves. Photographer Don Wood, Monga National Park south east of Baridwood



Branches with young seed cases. Photographer Jackie Miles



Branches with open seed cases. Photographer Jackie Miles



Bases of trees. Australian Plant Image Index, photographer Murray Fagg, Monga National Park south east of Baridwood



Trees. Australian Plant Image Index, photographer Murray Fagg, Monga National Park south east of Baridwood

## Common name

Eastern Leatherwood, Pinkwood, Plumwood, Stinkwood

## Family

Cunoniaceae

## Where found

**Rainforest** and gullies. Coast and ranges.

## Notes

Tree or **shrub** to 30 m high. Bark greyish brown, with longitudinal fissures and cracks, becoming more or less scaly. Branchlets brown- to white-hairy, sometimes becoming hairless. Leaves opposite each other, 5-17 cm long, **compound**, with 2-13 leaflets, or some leaves **simple**. Leaflets and simple leaves 1-7 cm long, 5-20 mm wide, leathery, dark green and glossy, or blotched whitish from dried resin, hairless or nearly hairless above, dull and white-hairy below, tips pointed to notched with a short **mucro**. Flowers with 4 white **petals** each 8-15 mm long. **Sepals** joined together and falling as a cap. Flowers in 1-5 flowered clusters. Flowers Jan.-Apr.

Family was Eucryphiaceae.

Vulnerable Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Eucryphia-moorei> (accessed 22 January, 2021)

---

*Author: Betty Wood.*



*Creative Commons Attribution 3.0 Australia (CC BY)*

*This identification key and fact sheets are available as a free mobile application:*



*Android edition*



*iOS edition*