Plants of South Eastern New South Wales



Flow ering stems. Photographer Russell Best, Wilsons Promontary, Vic



Flow ering stems. Photographer Don Wood, Eurobodalla Regional Botanic Gardens north of Moruya



Line drawings (as *Galium propinquum*). a. seeding branch; 'seed'. MMoir © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

A bedstraw

Family

Rubiaceae

Where found

Forest, woodland, and grassland. Widespread. No records from the Western Slopes in the area covered by this key.

Notes

Perennial herb, prostrate to sprawling, often extensively rhizomatous. Stems mostly to 0.4 m long, 0.5 mm in diameter, with slender angles, sparsely hairy to hairless. Leaves and leaf-like stipules in whorls of 4, 0.2–1.5 cm long, 1–5 mm wide, upper surface usually with scattered flattened clear triangular prickles, lower surface hairless or with a few flattened clear triangular prickles along the midrib, margins flat or narrowly curved down; tips pointed. Flowers 1–1.5 mm in diameter, greenish or greenish cream, with 4 petals joined together near their bases, in clusters of 1–6 flowers. Flowering: spring–summer. 'Seeds' 0.8–1.2 mm long, sparsely hairy to hairless, not or hardly wrinkled, dark brown.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Galium~leiocarpum (accessed 17 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





