Plants of South Eastern New South Wales



Flowers and leaves. Australian Flant Image Index, photographer Murray Fagg, near Batemans Bay



Flow ering stems. Photographer Don Wood, Bangalee Reserve near Nowra



Line drawings. a. stem-segment, flower and leaves; b. carpels. GRM Dashorst, State Herbariumof South Australia, © 2021 Royal Board, Cardens Board,

Common name

Twining guinea flower, Trailing guinea flower

Family

Dilleniaceae

Where found

Forest, sometimes in rocky areas. Coast and ranges.

Notes

Woody twiner with wiry stems to about 3 m long, sometimes trailing. Stems hairless to densely hairy with simple wart based hairs. Young stems purplish, older stems with scars that encircle the stems. Leaves alternating up the stems, 2.2–7 cm long, 9–30 mm wide, tips pointed to blunt, margins toothed, occasionally entire, surfaces hairy with simple hairs when young, usually becoming hairless with age, lower surface often purplish red, base blunt, with a distinct leaf stalk. Flowers with 5 yellow petals each 8–30 mm long. Stamens more than 30, surrounding the carpels. Carpels 3, hairless. Flowers single, at the bases of the leaves or at the ends of short shoots, stalks 0.3–19 mm long. Flowering: chiefly spring.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Hibbertia~dentata (accessed 19 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





