Plants of South Eastern New South Wales




Flow ering branches. Photographer Don Wood, Sassafras



Flow ering branches. Australian Plant Image Index, photographer Murray Fagg, Bobundara Nature Reserve east of Dalgety



Shrub. Australian Flant Image Index, photographer Murray Fagg, between Bungendore and Braidwood



Line drawings. e. flowering branch; f. seed case. SM Clarke, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Violet kunzea, Crimson kunzea

Family

Myrtaceae

Where found

Dry forest, woodland, and heath, and rocky slopes, often near seepage areas. Weedy after natural vegetation is cleared for grazing (personal communication from Roger Farrow, 25 January, 2021). Widespread. Rare in coastal areas.

Notes

Shrub to 2 m tall. Young stems hairy. Leaves aromatic when rubbed, crowded, alternating up the stems, mostly 0.1-0.4 cm long, 0.5-1 mm wide, tips more or less curved down, only the midvein prominent, surfaces at first hairy, becoming hairless with age, margins entire. Flowers with 5 pink to purple, rarely white, petals 1.5-2.5 mm long, in head-like clusters. Stamens 2-4 mm long. Flowers Spring to Summer.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Kunzea~parvifolia (accessed 25 January, 2021)

Author: Betty Wood.

 $This identification \ key\ and\ fact\ sheets\ are\ available\ as\ a\ free\ mobile\ application:$





