# **Plants of South Eastern New South Wales**



Flowers. Australian Flant Image Index, photographer Murray Fagg, Cooma



Flowering plant. Photographer Don Wood, The Pinnacle Reserve, Canberra, ACT



Plants. Photographers RG & FJ Richardson, Werribee, Vic

#### Common name

Prickly lettuce

### **Family**

Asteraceae

## Where found

Dry forest, woodland, grassy areas, roadsides, disturbed sites, and along streams. Widespread. Mostly in the Sydney area, tablelands, ACT, and Western Slopes.

# **Notes**

Introduced annual or biennial herb to to 3 m tall. Prickly, bristly, or hairless stems. Leaf margins and midrib spiny. Leaves alternating up the stems, usually held in a vertical plane, 3-20 cm long, 20-60 mm wide, glaucous, hairless, margins entire to dissected, basal lobes stem-clasping. Upper leaves becoming bract like below the flower heads. Flower heads rarely open. Open flower heads to about 20 mm in diameter, with 7-18 pale to bright yellow "petals' each 2.5–4 mm long. Flower heads below the 'petals' narrow cylindrical, 6-10 mm long. Flower heads in open clusters. Flowers Spring to Autumn.

- f. integrifolia: Lower leaves entire; mid-stem leaves entire or sometimes with a few shallow broad teeth; upper stem leaves and branch leaves entire
- f. serriola: Lower leaves usually deeply dissected, rarely margins entire; stem leaves lobed to dissected in the middle third of the stem

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Lactuca~serriola (accessed 22 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





