Plants of South Eastern New South Wales




Flow ering branches. Photographer Don Wood, Nerrigundah, w est of Bodalla



Trunks. Australian Pant Image Index, photographer A Lyne, Cotter Reserve, ACT



Young nuts and leaves. Australian Rant Image Index, photographer A Lyne Jamieson, Vic



Line drawings. f. branchlet, leaf, nut, seed. SM Clarke, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Slender tea-tree

Family

Myrtaceae

Where found

Dry forest, woodland, shrubland, stream banks, gullies, and rocky outcrops. Widespread.

Notes

Shrub or small tree to 4 m tall or more. Bark on smaller stems smooth, shedding in stringy strips, on larger stems rough, finely fissured. Young stems hairy, with a groove or a narrow flange near bases of the leaf stalks. Branches usually weep. Leaves aromatic when rubbed, alternating up the stems, 0.6–2.5 cm long, usually 2-5 mm wide, margins flat or curved down, surfaces silky to hairless; tips blunt or pointed, usually curved, with a minute, sharp point; bases narrowing to a short stalk. Flowers 6-11 mm in diameter, with 5 white petals, single or in pairs. Flowers Spring to Summer. Nuts shed early, usually 3-4 mm in diameter, surface silky to sparsely silky.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Leptospermum~brevipes (accessed 12 January, 2021)

Additional information in: Thompson, J. (8 December 1989), A revision of the genus *Leptospermum* (Myrtaceae). *Telopea* 3(3): 382-383, map 5-33

Author: Betty Wood.

6 ۲ This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)