Plants of South Eastern New South Wales



Flow ering branch and trunk. Photographer Don Wood, Yurammie State Forest, west of Merimbula




Young nut on branch. Photographer John Tann, Waterfall, Sydney



Open nut on branch. Photographer John Tann, Waterfall, Sydney



Trunk. Photographer John Tann, Waterfall, south of Sydney



Line drawings. a. branchlet, leaf, nut. SM Carke, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Paper-bark tea-tree, Slender tea-tree, Flaky-barked tea-tree

Family

Myrtaceae

Where found

Dry forest, heath, shrubland, and rocky areas. Widespread, mainly on the coast and ranges.

Notes

Shrub or small tree to 6 m tall. Bark rough, flaking in thin layers. Stems hairy. Leaves aromatic when rubbed, alternating up the stems, usually 1-2 cm long, 1-6 mm wide, silky, becoming hairless, tips rounded to pointed, with a blunt point; bases tapering or rounded, with a short stalk. Flowers 7-15 mm in diameter, with 5 white petals, single or in pairs. Flowers Spring to Summer. Nuts shed early, mostly 3-6 mm in diameter, surface hairy.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?

page=nswfl&lvl=sp&name=Leptospermum~trinervium (accessed 22 January, 2021)

Additional information in: Thompson, J. (8 December 1989), A revision of the genus *Leptospermum* (Myrtaceae). *Telopea* 3(3): 366-368, map 4-19

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)