# **Plants of South Eastern New South Wales**



Flowers and leaves. Photographer Don Wood, near Nerriga



Flowers and leaves. Australian Flant Image Index, photographer Murray Fagg, near Nowra

## Common name

Bird's-foot trefoil, Greater bird's foot trefoil

## **Family**

Fabaceae

#### Where found

Woodland, shrubland, moist grassy areas, along streams, and swamps. Widespread.

#### **Notes**

Introduced perennial herb to 0.8 m tall or prostrate. Stems hairless to sparsely hairy. Leaves alternating up the stems, compound, with 5 leaflets, each 0.4-2.5 cm long, 2-15 mm wide, tips pointed or more or less rounded, surfaces hairless to sparsely hairy, lower surface glaucous, the lower 2 leaflets usually smaller than the others. Flowers 10-12 mm long, pea shaped, with 5 petals, 2 joined together to form the keel, yellow to orange, often with reddish veins on the standard petal. Usually 8-14 flowers in a cluster. Flowers mainly Spring to Summer.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Lotus~uliginosus (accessed 24 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)