
Plants of South Eastern New South Wales



Flowers. Photographer Jackie Miles



Flowering stem. Photographer Jackie Mles



Line drawings. c. flowering branch; d. flower. BM Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Yellow loosestrife

Family

Primulaceae

Where found

Pasture, wetlands, and near streams. Wingecarribee Swamp, the Boro area near Braidwood, and the Bega River Valley.

Notes

Perennial herb to 2 m high, rhizomatous or stoloniferous. Dies back to the rootstock each autumn and resprouts next spring. Leaves in whorls of 3 or 4, or opposite each other, 3.5–9 cm long, 6–32 mm wide, becoming smaller towards the stem tip, green, paler beneath, covered with minute glandular hairs, dotted with orange to blackish glands. Flowers yellow, 9–15 mm in diameter, with 5 petals. Sepals with a line of orange to blackish glands on their outer surface near the margins. Flowers in many-flowered branched clusters.

Endangered NSW. Provisions of the NSW Biodiversity Conservation Act 2016 No 63 relating to the protection of protected plants generally also apply to plants that are a threatened species.

NSW Threatened Species profile: http://www.environment.nsw.gov.au/threatenedSpeciesApp/profile.aspx?id=10498 (accessed 7 January, 2021)

PlantNET description: https://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Lysimachia~vulgaris (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)