Plants of South Eastern New South Wales



Rowering stem Australian Rant Image Index, photographer Alexander Schmidt-Lebuhn, west of Miles, Qid

Common name

Native pear

Family

Apocynaceae

Where found

Woodland and shrubland. Coast, ranges, and the eastern edge of the tablelands, mostly north of Picton.

Notes

Climber with twining stems to 4 m high. Young stems hairy with fine hairs with colourless hairs, older stems and most leaves hairless. Leaves opposite each other, 2–12 cm long, 2–20 mm wide, rather thick, slightly fleshy, midvein prominent, channelled above, surfaces hairless or with scattered outgrowths below, bases sometimes cordate, glands minute, 1–5 at the base of the midvein. Flowers greenish or yellow, 3–6 mm in diameter, with a cup-shaped to bell-shaped tube, and 5 lobes, in 3–10 flowered umbels. Flowering: spring.

Rowering stem Australian Rant Image Index, photographer Alexander Schmidt-Lebuhn, west of Miles, Qid

Family was Asclepiadaceae.

Endangered population in the Bankstown, Blacktown, Camden, Campbelltown, Fairfield, Holroyd, Liverpool and Penrith local government areas.

Provisions of the Biodiversity Conservation Act 2016 No 63 relating to the protection of protected plants generally also apply to plants that are a threatened species or a part of a threatened ecological community.

NSW Threatened Species profile: http://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?id=10508 (accessed 7 January, 2021)

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=in&name=Marsdenia~viridiflora+subsp.~viridiflora (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





Android edition 🛛 🛃 iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)