Plants of South Eastern New South Wales



Rower (right) and unopened flower (left). Smithsonian Institution, National Museumof Natural History, Department of Botany, creator RA Howard, Hawaii



Young fruit and leaves. Australian Plant Image Index, photographer Murray Fagg, SW of Tweed Heads.

Common name

Fruit salad plant

Family

Araceae

Where found

Mainly where garden refuse has been dumped. Urban bushland and along streams. Sydney area and lower Blue Mountains south to near Kiama.

Notes

Introduced epiphyte, woody vine or perennial herb to 20 m tall, mostly to 1.5 m tall. Leaves alternating up the stems, 25–90 cm long, 250–750 mm broad, leathery, glossy, cordate, becoming lobed and holey in adult leaves. Individual flowers small. The spathe surrounds the cylindrical flower spike. Spike to 300 mm long. Fruit yellowish or green to purple when ripe, with a pungent smell. Edible only when fully ripe.

The description above is based on the article in Wikipedia and on the description in PlantNET of the family Araceae.

PlantNET description of the family Araceae: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=fm&name=Araceae (accessed 19 April 2021)

Wikipedia description: https://en.wikipedia.org/wiki/Monstera_deliciosa (accessed 19 April 2021)



Ripe fruit. Photographer B Navez, unknown place



Plant and flower. Photographer maliemociones, Mexico

Author: Betty Wood.

6 ۲ This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)