
Plants of South Eastern New South Wales



Flowers and leaves. Australian Plant Image Index, Photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ering carpet. Photographer Arthur Chapman, near Underbool, western Vic



Line drawings. a. Flowering branch; flower. MMbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Creeping Myoporum, Creeping Boobialla

Family

Scrophulariaceae

Where found

Clay soils, often on saline flats. Western Slopes.

Notes

Prostrate mat-forming shrub usually less than about 0.1 m tall. Fruit fleshy. Stems smooth to warty. Leaves alternating along the stems, clustered, 0.5-5.7 cm long, 1.5-8 mm wide, surfaces the same colour on each side, green to brown, hairless, often warty, margins entire, toothed, or warty, tips blunt or pointed. Flowers white or pink, with purple spots inside, 7–9 mm in diameter, with a bell-shaped tube and 5 lobes, the fronts of the lobes hairless or hairy. Flowers in 1-5 flowered clusters. Flowers most of the year. Fruit white or yellowish white, translucent, drying brown.

Family was Myoporaceae.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Myoporum~parvifolium (accessed 8 January, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$





