


Plants of South Eastern New South Wales



Flowers. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flower and young seed case. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flowering plant. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

Common name

Bower vine, Bower of beauty

Family

Bignoniaceae

Where found

Garden **escape**. Coastal, mainly in the Sydney area.

Naturally grows in **rainforest** north from west of Port Macquarie.

Notes

Introduced tall woody climber. Older stems longitudinally grooved. Stems hairless. Leaves mostly opposite each other, sometimes in **whorls** of 3, 12–20 cm long, **compound**, with 5–9, rarely 3, leaflets, each 4.5–8 cm long, 15–40 mm wide, tips pointed, margins **entire**, surfaces hairless. Flowers white or pink with red markings in the throat, 40–60 mm long, tubular, the tube 40–60 mm long, 10–20 mm in diameter, with 5 **spreading** lobes each 20–30 mm long. Flower clusters 6–12 cm long. Flowering: September–March. Seeds with a membranous **wing**.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pandorea~jasminoides> (accessed 10 April 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)