Plants of South Eastern New South Wales



Flowers and leaves. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ering stem and leaves. Photographer Richard Hartland, Tallageira, Vic



Flowers and leaves. Photographer David Ptts, © 2021 Ptts, David. Published in VIOFLORA

Common name

Coast Stork's-bill

Family

Geraniaceae

Where found

Usually on sandy soils. Widespread.

Notes

Perennial herb to about 0.4 m tall. Stems hairy, sticky. Leaves opposite each other, to 6 cm long, to 60 mm wide, oval with cordate bases or round, entire to shallowly 5–7-lobed, surfaces usually sparsely hairy with coarse hairs or becoming hairless, margins scalloped or bluntly toothed. Flowers usually deep pink, with 5 petals each 5–6 mm long, and 3-5 stamens that produce pollen. Sepals 3–4 mm long, narrow, tips pointed, with sparse, long (about 1 mm or more) spreading hairs, sometimes mixed with much shorter erect hairs, sepal spur 1–3.5 mm long. Flowers in 2–7-flowered clusters. Flowers Spring to Autumn.

Data deficient Vic.

VICFLORA description: https://vicflora.rbg.vic.gov.au/flora/taxon/49ec3198-acd7-424d-b967-5e30ea69582c (accessed 8 March 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)