
# **Plants of South Eastern New South Wales**



Flow er head. Photographer Richard Hartland, Bogong High Plains, Vic



Rowering plants. Photographer Russell Best, Bogong High Plains, Vic



Line drawings. b. plant; leaf upper surface detail. E Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

#### Common name

**Broad Plantain** 

### **Family**

Plantaginaceae

## Where found

Alpine and subalpine **grassland** and **herbfields**. Western edge of the ACT and the mountains to the west. Kosciuszko National Park and nearby. Tablelands and western edge of the ranges south of Captains Flat.

#### Notes

Perennial herb to 0.3 m high, with fleshy adventitious roots. Leaves fleshy. Flower stalks densely hairy with rusty hairs. Leaves in a basal rosette, 3–11 cm long, 10–45 mm wide, hairy, more densely so on the lower surface, with 3-9 longitudinal veins, margins entire or with a few small teeth, tips blunt. Axils with tufts of long golden-brown hairs. Flowers tubular, the tube 1.5–2.5 mm long, with 4 lobes 0.5–1 mm long, spreading or turned down. Sepals 1.5–2.5 mm long. Sepal-like bracts just beneath the flowers 2–4 mm long. Flower spikes 10 to many-flowered, narrow cylindrical, dense, 20–120 mm long at the seeding stage. Flowering: December–March

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Plantago~euryphylla (accessed 25 January, 2021) Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)