Plants of South Eastern New South Wales



Row ering stems. Australian Rant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Leafy stems with spent flowers. Photographe Jackie Mles



Row ering shrub. Australian Rant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

Common name

Veined pomaderris

Family

Rhamnaceae

Where found

Open **forest**, **shrubland**, rocky sites, and near streams. ACT, Tablelands in Tallaganda State Forest. Ranges south from Deua National Park. Coastal south of Pambula.

Notes

Shrub to 4 m high. New growth and young stems covered with fawn to golden or rusty simple and stellate hairs (needs a hand lens or a macro app on your phone/tablet to see). Leaves alternating up the stems, 2–5 cm long, 10–35 mm wide, upper surface hairless; lower surface densely hairy with whitish or pale to golden hairs, often obscuring the underlying stellate hairs. Hairs simple and mostly lying in the same direction; lateral veins pale brown to rusty, looping to the inside and not reaching the leaf margins, tips mostly blunt, margins flat or weakly wavy. Flowers with 5 white to yellow sepals 1.2–1.8 mm long, falling early, and 0 petals, in large many-flowered branched clusters 30–50 mm long.

All native plants on unleased land in the ACT are protected.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pomaderris~costata (accessed 2 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:


é

iOS edition

Android edition



Creative Commons Attribution 3.0 Australia (CC BY)



Line drawings. f. flower; leaf outline; leaf lower surface detail. A. Barley, National Herbarium of Victoria, \odot 2021 Royal Botanic Gardens Board