Plants of South Eastern New South Wales



Flower head with leafy bracts. Australian Flant Image Index, photographer Murray Fagg, Kosciuszko National Park



Flow ering stems. Photographer Don Wood, Mungerarie State Forest west of Moruya



Line drawing. USDA NRCS PLANTS Database: Britton, N.L., and A. Brown. 1913. *An illustrated flora* of the northern United States, Canada and the British Possessions. Vol. 3: 115.

Common name

Self-heal, Heal All

Family

Lamiaceae

Where found

Woodland, pastures, disturbed areas, roadsides, moist sites, and stream banks. Widespread.

Notes

Introduced perennial herb to 0.5 m high, often sprawling to prostrate, often rooting at the lower nodes. Stems 4-angular, almost hairless except for a few hairs mainly near the nodes. Leaves aromatic when rubbed, opposite each other, 1.5–6 cm long, 7–25 mm wide, surfaces with scattered coarse hairs, tips blunt to somewhat pointed, margin entire to obscurely toothed. Flowers deep purpleblue to mauve or pink, 8–12 mm long, tubular, with 5 lobes forming 2 lips. Flowers in a dense, 1–6 cm long cluster. Flowering: November–April.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Prunella~vulgaris (accessed 3 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





